



上海中醫藥大學

Shanghai University of Traditional Chinese Medicine


2014

GB/T7713-1987

GB/T 7714-2015

GB/T7713.1-2006

1.1	1
1.1.1	1
1.1.2	1
1.1.3	1
1.1.4	1
1.1.5	1
1.1.6	1
1.2	2
1.3	2
1.4	2
1.5	3
1.6	3
1.6.1	4
1.6.2	4
1.7	4
1.7.1	5
1.7.2	5
1.7.3	5
1.7.4	6
1.8	8
1.8.1	8
1.8.2	11
1.8.3	12
1.9	12
1.10	12
21	14
22	14



--	--	--	--

•

•

•

•

•

•

表 3 非浸润性膀胱癌不同中医证型患者生存质量评分比较($\bar{x}\pm s$, 分)

证型	n	乏力	疼痛	呼吸困难	失眠	食欲丧失	便秘	排尿症状	肠道症状
瘀毒蕴结型	144	64.4±22.7	69.7±23.4*	5.9±15.3	44.3±24.5	40.9±27.6	55.5±18.3*	52.9±19.6*	39.5±18.8*
湿热下注型	149	64.3±23.5	57.6±21.3#	5.8±15.8	45.1±23.4	42.7±25.1	23.4±8.1	68.8±16.7#	58.4±18.3#
脾肾两虚型	18	67.5±22.8	52.8±20.7**	6.4±15.2	45.6±22.9	57.5±23.4*#	47.9±22.5*#	46.2±16.5**	47.1±17.9**
肝肾阴虚型	44	63.4±24.9	55.3±22.5	6.2±15.8	47.0±20.6	48.4±23.1	59.9±20.8	50.7±19.4	38.6±19.2
无证可辨/其他证型	31	43.0±22.8*	40.4±23.4*	6.1±15.8	43.3±24.3	41.1±23.5	39.3±24.7*	41.5±18.6*	36.7±15.0*

证型	n	经济困难	躯体功能	认知功能	情绪功能	社会功能	总体健康状况	对将来担心	生活质量
瘀毒蕴结型	144	57.6±31.5	60.1±18.4	78.3±20.1	41.7±24.6	35.4±22.7	21.6±14.3	51.6±10.3	7.5±16.6
湿热下注型	149	53.4±27.3	57.5±18.0	78.6±21.7	39.8±17.4	39.1±19.9	23.4±8.1	50.7±12.1	6.8±13.4
脾肾两虚型	18	52.1±28.8	58.2±18.1	79.4±21.3	38.4±17.2	32.5±21.0*#	19.5±8.5*	51.5±11.8	6.6±16.2
肝肾阴虚型	44	58.4±32.3	56.0±19.5	79.1±21.2	42.2±24.5	36.6±23.8	24.9±13.4	50.4±11.6	4.9±11.5*#
无证可辨/其他证型	31	56.4±31.3	65.3±18.2*#	77.5±24.8	47.5±15.3*#	40.8±24.3	40.3±6.5*#	52.9±15.3	6.8±15.8

注:与湿热下注型比较,* $P<0.05$;与瘀毒蕴结型比较,# $P<0.05$ 。

Table 2 Treatment and the prognosis of recurrence breast cancer patients

Index	TNBC	HER2+	HER2-	P
Surgery type				
Total mastectomy	382(86.15)	230(85.74)		
Distant recurrence				
>24	187(41.10)	1590(59.20)	51.867	<0.001
Type of recurrence				
Distant metastasis	357(78.46)	2112(78.63)		
Local recurrence	89(19.56)	486(18.09)	2.582	0.275
Unknown	9(1.98)	88(3.28)		
Liver metastasis				
Yes	71(15.60)	562(20.92)		
No	384(84.40)	2117(78.82)	8.169	0.017
Unknown	0	7(0.26)		
Lung metastasis				
Yes	161(35.29)	707(26.4)		
No	294(64.62)	1885(70.18)	7.262	0.026
Unknown	0	7(0.26)		
Bone metastasis				
Yes	129(28.35)	1041(38.76)		
No	326(71.65)	1638(60.98)	19.573	<0.001
Unknown	0	7(0.26)		
Brain metastases				
Yes	17(3.74)	84(3.13)		
No	438(96.26)	2595(96.61)	1.639	0.441
Unknown	0	7(0.26)		

表 1 两组患者临床资料比较 [例 (%)] $\bar{x}\pm s$

临床资料	异丙托溴铵组 (96例)	安慰剂组 (94例)	整体人群 (190例)
性别			
男	83 (86.5)	85 (90.4)	168 (88.4)
女	13 (13.5)	9 (9.6)	22 (11.6)
年龄(岁)	63.4±6.5	62.4±6.4	62.9±6.5
年龄≥60岁	67 (69.8)	65 (69.1)	132 (69.5)
急性支气管扩张剂反应性			
是	13 (13.5)	13 (13.8)	26 (13.7)
否	83 (86.5)	81 (86.2)	164 (86.3)
身高 (cm)	165.90±7.30	168.60±7.90	167.20±7.70
体重 (kg)	61.80±11.58	64.07±11.66	62.93±11.65
体重指数 (kg/m ²)	22.38±3.42	22.50±3.51	22.44±3.46
吸烟状况			
从不吸烟	15 (15.6)	6 (6.4)	21 (11.1)
以前吸烟	55 (57.3)	54 (57.4)	109 (57.4)
目前吸烟	26 (27.1)	34 (36.2)	60 (31.6)
饮酒史			
不饮酒者	60 (62.5)	54 (57.4)	114 (60.0)
饮酒者	36 (37.5)	40 (42.6)	76 (40.0)



示例如下：

-

示例如下：

-

示例如下：

示例如下：




示例如下：



示例如下：







-
-
-
-
-
-
-
-
-
-
-